

Leaf-gall thrips of *Ficus*—*Gynaikothrips ficorum* (Marchal) and *Gynaikothrips uzeli* (Zimmerman) (Insecta: Thysanoptera: Phlaeothripidae)¹

H. A. Denmark, Thomas R. Fasulo, and Joseph E. Funderburk²

Introduction

Leaf-gall thrips in the genus *Gynaikothrips* originating in southeast Asia infest *Ficus*. This is not a recent introduction. Specimens collected in Florida in 1887 are in the U.S. National Museum (Denmark, 1967). Watson (1923) listed the genus in his key to thrips of North America. Two species, recognized as distinct by Priesner (1939) based only on differences in the length of the pronotal setae, are established in Florida: *Gynaikothrips ficorum* (Marchal) and *Gynaikothrips uzeli* (Zimmerman). However, whether or not they are distinct species is uncertain. Populations in Southeast Asia exhibit considerable variation in the lengths of the pronotal setae, and Mound et al. (1996) suggested that *Gynaikothrips ficorum* is probably a form of *Gynaikothrips uzeli* that has been spread by the horticultural trade. *Gynaikothrips ficorum* and *Gynaikothrips uzeli* are commonly called Cuban laurel thrips and weeping fig thrips, respectively.

Distribution

While most *Gynaikothrips* spp. are of Asian origin, species in this genus have been described from Africa. *Gynaikothrips ficorum* is pantropical, appearing wherever *Ficus microcarpa* L. f. is planted (Denmark, 1967). It is recorded



Figure 1. Several life stages of the Cuban laurel thrips, *Gynaikothrips ficorum* (Marchal). Black forms are adults, yellow forms are immatures and white, capsule-shaped forms are eggs.
Credits: Paul M. Choate, University of Florida

from Algeria, the Canary Islands, Colombia, Cuba, Dominican Republic, Guam, Taiwan, Ecuador, India, Java, Mexico, Nassau (Bahamas), Nicaragua, Israel, Palestine, Panama, Puerto Rico, El Salvador, Thailand, Spain, Sicily, and the United States. In the USA, it is recorded from California, Florida, Hawaii, and Texas. Once it became established in

1. This document is EENY-324 (IN599) (originally published as DPI Entomology Circular 59), one of a series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Published: August 2004. Revised: April 2014.. Reviewed: March 2011. This document is also available on Featured Creatures website at <http://entomology.ifas.ufl.edu/creatures>. Please visit the EDIS website at <http://edis.ifas.ufl.edu>.
2. H. A. Denmark, Florida Department of Agriculture and Consumer Services, Division of Plant Industry; and Thomas R. Fasulo and Joseph E. Funderburk, University of Florida.

Florida, *Gynaikothrips uzeli* was rapidly spread on *Ficus benjamina* L. originating from nurseries in southern Florida (Held et al., 2005). It has been reported from 10 states in the continental USA: Hawaii, Puerto Rico, Trinidad, Mexico, Colombia, Argentina, and Costa Rica (Cabrera-Asencio et al., 2008; Held and Boyd, 2008a; de Borbon and Agostini, 2011).

Description

The life cycle takes about 30 days (Arthurs et al., 2011). Adults migrate to terminal leaves and establish folded-leaf galls within 2 to 3 days of infestation. Mating, egg laying, and a complete generation develop within a single gall. Adults exit galls within a few days of emergence, and migrate to new terminal leaves, either on the same or different terminal stem, to begin a new generation.

Egg: The egg is cylindrical with rounded ends, smooth, and is a translucent white.

Larva: There are two larval instars. The first instar is small and translucent white. The second instar is larger, light yellow and the posterior tube is dark and is held pointing up. The second instar is similar to the adult in size and shape. Both the first and second instars have red eyes.

Prepupa and Pupa: The prepupa has wing buds that are externally visible. The pupa has longer wing buds and the antennae are folded back over the head.

Adult: The adult varies from about 2.6 mm to 3.6 mm in length and is dark yellowish-brown to black. With the exception of the legs and last abdominal segment, dorsal striations are shown on the adult. Breeding is continuous. Although the adult moves or flies rapidly when disturbed,



Figure 2. Pupa of the Cuban laurel thrips, *Gynaikothrips ficorum* (Marchal). The pupae are easily distinguished from prepupae by their longer wing buds and the antennae folded back over the head. Credits: Paul M. Choate, University of Florida

it remains on or close to the leaves most of the time. They are also active flyers on hot days. Canizo (1945) and de Borbon and Agostini (2011) give detailed measurements of *Gynaikothrips ficorum* and *Gynaikothrips uzeli*, respectively.

Hosts

Chinese banyan, *Ficus microcarpa* L. (Urticales: Moraceae) is host for *Gynaikothrips ficorum* (Mound et al., 1996). Weeping fig, *Ficus benjamina* L., is host for *Gynaikothrips uzeli* (Arthurs et al., 2011). Wolcott (1953) says that most species of *Ficus* are largely or entirely immune to infestations, but when the thrips become abundant, they will feed on certain non-host plant species. Cuban laurel thrips are an occasional pest of orchids (Nixon, 1999).



Figure 3. Adult Cuban laurel thrips, *Gynaikothrips ficorum* (Marchal). Notice the empty egg sac and shed exuviae of immatures. Credits: Paul M. Choate, University of Florida

Refer to Hoddle et al. (2012) for more information on *Gynaikothrips*, including a key to the species of adult thrips. [Key to eleven species of thrips.](#)

Damage and Economic Importance

Adult thrips use their sucking mouthparts to feed on the tender, light-green leaves, causing sunken purplish-red spots. Feeding results in a specific, directed growth reaction that causes the leaf to roll, or the leaf may fold along the midrib. The curled leaf becomes tough as it yellows, sometimes dropping during rainy or windy weather. Leaves eventually drop from the plant prematurely. Infested trees will not be killed, but the ornamental value of the plant is reduced markedly. The thrips rarely are annoying but occasionally may bite people. Some countries restrict the importation of plant material intercepted at ports-of-entry

when they are infested with *Gynaikothrips ficorum* or *Gynaikothrips uzeli*.

Some exotic *Ficus* species are listed as invasive in Florida. Bennett (1995) suggested that *Gynaikothrips* potentially may serve as biological control agents of these unwanted invaders. Refer to Mound (2013) for more information on thrips and galls, including leaf-gall thrips of *Ficus*.

Management

Pest status, the intensity of management, and the types of tactics differ in landscape and commercial production. Considerable injury is generally tolerable on landscape plantings of *Ficus* whereas detectable populations or plant injury on nursery plants is generally intolerable.



Figure 4. Injury on the leaves of *Ficus* due to feeding by *Gynaikothrips ficorum* (Marchal).

Credits: Paul M. Choate, University of Florida

Biological control: Biological control: *Montandoniola confusa* Streito and Matocq (Hemiptera: Anthocoridae) is the most widespread and important natural enemy in the southeastern US. It was found in Florida in 1990 (Dobbs and Boyd, 2006), and incorrectly identified as *Montandoniola moraguesi* (Bennett, 1995; Pluot-Sigwalt et al., 2009). The arrival of this species originally from the Philippines appears to be completely serendipitous. It is a very effective predator of *Gynaikothrips*, reducing populations more than 95%, and leaf galls by 77% (Arthurs et al., 2011); therefore, relying on this natural enemy as a biological control is the best option in the landscape. The native anthocorid *Orius insidiosus* (Say) is an excellent predator of many species of thrips, but it is not effective in reducing the *Gynaikothrips* in the leafgalls. Another thrips described from India, *Androthrips ramachandrai* Karny (Thysanoptera: Phlaeothripidae), has become established in the Americas, from the United States to Argentina. This thrips inhabits the leaf-galls and is an effective predator of *Gynaikothrips*

(de Melo et al., 2013). Refer to Boyd and Held (2006) and de Borbon and Agostini (2011) for more information on *Androthrips ramachandrai*.

Other potential biological controls include green lacewing larvae, bigeyed bugs, damsel bugs, ladybird beetles, parasitic wasps, predatory mites, and *Verticillium* (a fungal pathogen) (Padrick, 2004; Held et al., 2005; Dobbs and Boyd, 2006).

Cultural control: Replacing *Ficus microcarpa* and *Ficus benjamina* with a resistant species of *Ficus* probably would be the best and most lasting control of this pest. Because both species only attack the tender new foliage, the new growth can be pruned to remove injured leaves and to suppress the thrips population.



Figure 5. Larva of the minute pirate bug, *Orius insidiosus* (Say).

Credits: James F. Price, University of Florida

Chemical control: Insecticides are available that are effective against *Gynaikothrips* on *Ficus* (Held and Boyd, 2008b). Systemic insecticides provide better control of thrips in galls. Broad-spectrum insecticides are detrimental to natural enemies that have the potential to naturally maintain populations of *Gynaikothrips* at relatively low numbers in the landscape (Arthurs et al., 2011). Insecticides are frequently the only option in nurseries considering the very low level of injury that is acceptable in domestic and international markets. University of Florida trials have shown that soil drench with certain systemic insecticides is effective, whereas foliar applications are not. Contact your county extension office for specific insecticide recommendations.



Figure 6. Adult and immature of *Montandoniola moraguesi* (Puton), a predator of the Cuban laurel thrips, *Gynaikothrips ficorum* (Marchal), shown here with an alternate host *Gynaikothrips uzeli* (Zimmerman 1900).

Credits: L.J. Buss, University of Florida

Selected References

Arthurs S, Chen J, Dogramaci M, Ali AD, Mannion C. 2011. Evaluation of *Montandoniola confusa* Streito and *Matocq* sp. nov. and *Orius insidiosus* Say (Heteroptera: Anthocoridae), for control of *Gynaikothrips uzeli* Zimmerman (Thysanoptera: Phlaeothripidae) on *Ficus benjamina*. *Biological Control* 57: 202-207.

Bennett FD. 1995. *Montandoniola moraguesi* (Hemiptera: Anthocoridae), a new immigrant to Florida: Friend or Foe? *Vedalia* 2: 3-6.

Boyd DW, Held DW. 2006. *Androthrips ramachandria* (Thysanoptera: Phlaeothripidae): an introduced thrips in the United States. *Florida Entomologist* 89: 455-458.

Cabrera-Asencio L, Ramirez A, Saez L, Velez AL. 2008. *Gynaikothrips uzeli* Zimmerman (Thysanoptera: Phlaeothripidae) y *Montandoniola moraguezi* Puton (Hemiptera: Anthocoridae): Nuevos records para Puerto Rico. *Journal of Agriculture of the University of Puerto Rico* 92: 111-113.

Canizo JD. 1945. Redescrición de *Gynaikothrips ficorum* (Marchal) concepto actual del género *Gynaikothrips* Zimmerman, nuevo para la fauna continental europea (Thysanoptera, Phlaeothripidae). *Eos*. 21: 123-156.

de Borbon CM, Agostini JP. 2011. *Gynaikothrips uzeli* (Zimmermann) and *Androthrips ramachandria* Karny

(Thysanoptera, Phlaeothripidae), first records for Argentina. *Revista de la Facultad de Ciencias Agrarias, Universidad Nacional de Cuyo* 43 (1): 253-260. (17 March 2014)

de Melo FS, Cavalleri A, de Souza Mendonca, Jr M. 2013. Predation of *Gynaikothrips uzeli* (Thysanoptera: Phlaeothripidae) by *Androthrips ramachandria* (Thysanoptera: Phlaeothripidae). *Florida Entomologist* 96: 859-863.

Denmark HA. 1967. Cuban laurel thrips, *Gynaikothrips ficorum*, in Florida. Florida Department of Agriculture. *Entomology Circular* 59, 1-2.

Dobbs TT, Boyd DW. 2006. Status and distribution of *Montandoniola moraguesi* (Hemiptera: Anthocoridae) in the continental United States. *Florida Entomologist* 89: 41-46.

Held DW, Boyd D. 2008a. New records of *Gynaikothrips uzeli* (Zimmerman) (Thysanoptera: Phlaeothripidae) on *Ficus benjamina* in Texas and O'ahu, Hawaii. USA. *Pan-Pacific Entomologist* 84: 77-80.

Held DW, Boyd D. 2008b. Evaluation of sticky traps and insecticides to prevent gall induction by *Gynaikothrips uzeli* (Zimmerman) (Thysanoptera: Phlaeothripidae) on *Ficus benjamina*. *Pest Management Science* 64: 133-140.

Held DW, Boyd D, Lockley T, Edwards GB. 2005. *Gynaikothrips uzeli* (Thysanoptera: Phlaeothripidae) in the southeastern United States: distribution and review of biology. *Florida Entomologist* 88: 538-540.

Hoddle MS, Mound LA, Paris DL. 2012. *Thrips of California*. CBIT Publishing, Queensland. (14 March 2014)

Marchal P. 1908. Sur une nouvelle espèce de thrips (Thysanopt.) nuisible aux *Ficus* en Algérie. *Bulletin de la Société Entomologique de France* No. 14: 251-253.

Mound LA. 2013. *Thrips and galls*. CSIRO Ecosystem Services, Canberra, Australia. (15 March 2014)

Mound LA, Wang C-L, Okajima S. 1996. Observations in Taiwan on the identity of the Cuban laurel thrips (Thysanoptera, Phlaeothripidae). *Journal of the New York Entomological Society* 103: 185-190.

Nixon P, et al. (1999). Orchid pests and their management. Home, Yard & Garden Pest Newsletter. Available at <http://www.ag.uiuc.edu/cespubs/hyg/html/pdf/199919.pdf> [accessed 20 March 2014].

Padrick RC. 2004. Tips for Cuban laurel thrips. *Ornamental Outlook* 12(7): 40.

Pluot-Sigwalt D, Streito JC, Matocq A. 2009. Is *Montandoniola moraguesi* (Puton, 1896) a mixture of different species? (Hemiptera: Heteroptera: Anthocoridae). *Zootaxa* 2208: 25-43.

Priesner H. 1939. Zur Kenntnis der Gattung *Gynaikothrips* Zimm. (Thysanoptera). *Mitteilungen der Muenchner Entomologischen Gesellschaft* 29: 475-487.

Watson JR. 1923. Synopsis and catalog of the Thysanoptera of North America. University of Florida Agricultural Experiment Station Bulletin 168: 100 p.

Wolcott GN. 1953. Control of the laurel thrips, *Gynaikothrips ficorum* (Marchal). *Journal of Agriculture of the University of Puerto Rico* 37: 234-240.